

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Indar<sup>®</sup> 75WSP

EPA Reg. No. 62719-421

### Disease Control in Blueberry

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Indar<sup>®</sup> 75WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Indar 75WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Indar 75WSP.

#### Directions for Use

Refer to product label for Indar 75WSP for Mixing and Application Instructions.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
alternaria ( <i>Alternaria</i> spp.) anthracnose ( <i>Colletotrichum gloeosporioides</i> ) leaf spot and blotch ( <i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) mummy berry disease ( <i>Monilinia vaccinii-corymbosi</i> ) phomopsis twig blight and fruit rot ( <i>Phomopsis vaccinii</i> ) powdery mildew ( <i>Microsphaera vaccinii</i> ) rusts ( <i>Pucciniastrum</i> spp.)	2 (0.094 lb active)	For mummy berry and phomopsis twig blight, begin applications at early green tip and make subsequent applications at intervals of 8 to 14 days. For control of powdery mildew and other diseases listed, begin applications prior to or at onset of disease development and repeat at 8- to 14-day intervals. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> <li>• Do not make more than 4 applications or apply more than 8 oz of Indar 75WSP (0.38 lb active) per acre per year.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> <li>• Do not graze livestock in treated fields or offer treated material as a livestock feed item.</li> </ul>

#### Resistance Management

Indar 75WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Indar 75WSP should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

#### Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R845-052

EPA accepted: \_\_/\_\_/\_\_

Initial printing.

# Indar<sup>®</sup> 75WSP

EPA Reg. No. 62719-421

## Registration Notes:

**Proposed Section 3 supplemental labeling.**

<sup>®</sup>Trademark of Dow AgroSciences LLC